



Treat the  
pain & rising  
body  
temperature  
with  
**comprehensive**  
choice for  
complete  
**relief**

# Nowcef

Cefixime (U.S.P.)

400mg Capsules,  
100 mg/5 ml  
&  
200 mg/5 ml  
Oral Suspension

*Nitro Fresh*  
  
**Strawberry**  
Flavoured

 **NAWAN**  
LABORATORIES (PVT) LTD.



# Nowcef

Cefixime (U.S.P.)

400mg Capsules,  
100 mg/5 ml  
&  
200 mg/5 ml  
Oral Suspension

Nitro Fresh  
  
Strawberry  
Flavoured



ناؤسیف  
(سیفکزم - یو ایس پی)

**Nowcef** IS EFFECTIVE IN THE TREATMENT OF TYPHOID FEVER IN CHILDREN WITH CLINICAL CURE RATE OF 93%.<sup>1</sup>

**Nowcef** IS BENEFICIAL IN ACUTE GASTROENTERITIS IN CHILDREN WITH MUCOID OR BLOODY DIARRHEA.<sup>2</sup>

**Nowcef** HAS HIGHER MICROBIOLOGICAL EFFICACY AND LESS SIDE EFFECTS IN THE TREATMENT OF ACUTE CYSTITIS.<sup>3</sup>

**Nowcef** PROVIDES SUCCESSFUL CLINICAL RESPONSE AGAINST COMMUNITY ACQUIRED INFECTIONS SUCH AS: AOM & PNEUMONIA.<sup>4</sup>

**Nowcef** IS THE RECOMMENDED ORAL TREATMENT IN GONORRHEA.<sup>5</sup>

**Dosage: Adults:** The recommended dose of cefixime is 400 mg daily. For the treatment of uncomplicated cervical/urethral gonococcal infections, a single oral dose of 400 mg is recommended. The capsule may be administered without regard to food.

In the treatment of infections due to *Streptococcus pyogenes*, a therapeutic dosage of cefixime should be administered for at least 10 days.

**Pediatric Patients (6 months or older):** The recommended dose is 8 mg/kg/day of the suspension. This may be administered as a single daily dose or may be given in two divided doses, as 4 mg/kg every 12 hours.

**Storage Condition:** Store below 30°C in a dark and dry place away from children.

MANUFACTURED BY:  
 **NAWAN**  
LABORATORIES (PVT) LTD. | 136, Sector 15, Korangi Industrial Area, Karachi-Pakistan.  
Tel+ 92 21 35065835-7  
Web: www.nawanlab.com

Reference:  
1. Begum, B. et al. "Comparison between Azithromycin and Cefixime in the treatment of Typhoid Fever in Children." *Mymensingh medical journal: MMJ* 23.3 (2014): 441-446.  
2. Lee, H.-Y., et al. "Oral Cefixime is beneficial in acute uncomplicated *Salmonella* gastroenteritis in children with mucoid or bloody diarrhea-Do indication of antibiotic therapy need reassessment?" *International Journal of Infectious Diseases* 14 (2010): e196-e197.  
3. Collins, W.J., et al. "Efficacy and safety of cefixime and ciprofloxacin in acute cystitis (a multicenter randomized trial)." *Urologia (Moscow, Russia)* 1999; 1 (2010): 13-16.  
4. Dreshaj, Sh. et al. "Clinical role of Cefixime in community-acquired infections." *Priloz/Makedonska akademija na naukite i umetnostite, Odselenie za bioslozhi medinski nauki- Contributions/Macedonian Academy of Sciences and Arts, Section of Biological and Medical Science* 32.2 (2011):143-155.  
5. Donders, Glibber GG. "Treatment of sexually transmitted bacterial disease in pregnant women." *Drug* 59.3 (2000): 477-485.