

Nawacin

Injection

50

(Oxytetracycline)



Nawacin

Injection
(Oxytetracycline)

50

COMPOSITION: Each 100ml contains:
Oxytetracycline (as HCl) 5 gm
(USP Specification)

TARGET SPECIES:

Cattle, Sheep, Goats, Dogs and Cats.

INDICATIONS:**Cattle:** actinobacillosis, shot, pneumonia, pasteurellosis, mastitis, metritis, colibacillosis. **Sheep and goats:** puerperal infections, mastitis, pneumonia. **Dogs and cats:** respiratory, urogenital and gastrointestinal infections, septicemia.

DOSAGE AND ROUTE OF ADMINISTRATION:

Parenteral administration: deep intramuscular or intravenous injection. Oxytetracycline is administered in doses of 5 – 10 mg/kg body weight. This corresponds to 1 – 2 ml per 10 kg body weight daily. In acute cases, it is advisable to start treatment with intravenous administration of 2 ml 10 kg body weight followed by intramuscular administration of 1 ml / 10 kg body weight daily. Do not administer intravenously to dogs and cats.

WITHDRAWAL PERIODS:

Meat and offal: 22 days. Milk: 7 days.
Dogs & Cats: Not Applicable.

INSTRUCTIONS:

Store below 25°C.
Protect from heat & light.
Keep out of reach of children.

