

Resistance to drenches is getting worse...

Use

Pamezan

D R E N C H

Levamisole, Oxytoclozanide, Cobalt & Selenium



for total



performance

 **NAWAN**
LABORATORIES (PVT.) LTD

EACH 100 ML CONTAINS:

LEVAMISOLE HCl B.P	1.5 gm
OXYCLOZANIDE B.P. (VET)	3.0 gm
COBALT SULPHATE B.P.	0.382 gm
SODIUM SELENITE B.P.	0.035 gm

Pamezan

D R E N C H

Levamisole, Oxyclozanide, Cobalt & Selenium

INDICATIONS:

Pamezan is a highly effective combination of potent and powerful anthelmintics (levamisole and oxyclozanide).

Pamezan controls and eliminates round worms, Tape worms, Lung worms and Fasciola species (Liver flukes) in immature and mature stages, including bile ducts fascioliasis in the cattle, cow, sheep and goats.

Pamezan has immunomodulating properties.

Pamezan strengthens the immunity against disease.

Pamezan having **cobalt** and **selenium** as nutritional supplement ensures healthy growth of skin, hair, wool and corrects the deficiencies in small and large animals.

DOSAGE:

Sheep & Goats

Weight (kg):	15	25	35	45	upto 50
Dose (ml) :	5	10	15	20	25

Cows & Buffaloes

Weight (kg):	75	125	225	300	upto 300
Dose (ml) :	25	50	100	125	150

Route of Administration : Oral

OPERATORS WARNING: Direct contact with skin should be kept to a minimum. Wash hands after use.

CONTRAINDICATION: Not to be used in animals with known hypersensitivity to the levamisole and oxyclozanide.

STORAGE CONDITIONS:

Store in cool dry place below 25° C, Protected from light
Keep out of the reach of children.

WITHDRAWAL PERIOD:

Meat 28 days
Milk 5 days

PHARMACOLOGY:

- **Pamezan** affect the transmission of nerve impulses in the worm resulting in paralysis and expulsion of the nematodes.
- Broad Spectrum Anthelmintic
- Rapid Absorption and Excretion
- Highly Soluble

WARNINGS AND PRECAUTIONS:

1). Shake well before use. 2). Not to be diluted or mixed with other products. **3).** Muscles Tremor and increased salivation can occur if dosage is exceeded. **4).** Pregnant and lactating animals should be treated with care. **5).** Care should be taken to estimate accurately the bodyweight of animals to be treated.

Packing: Available in pack of 150 ml and 1 litre

