



QUINAMIN

Quinapyramine Sulphate
Quinapyramine Chloride

For Treatment and Prophylaxis
of Trypanosomiasis

QUINAMIN

Quinapyramine Sulphate
Quinapyramine Chloride



For Treatment and Prophylaxis of Trypanosomiasis

Composition:

Each vial contains:

Quinapyramine Sulphate B. Vet. C. 1.5 gm
Quinapyramine Chloride B. Vet. C. 1.0 gm

Indications: Quinamin is indicated for the treatment and prophylaxis of Trypanosomiasis caused by species of Trypanosomes-T.evansi in camels.

Dosage:

Add 15 ml sterile water for injection and shake well to make suspension for immediate use.

The recommended dose is 0.025 ml of Quinamin injection per kg body weight.

Do not exceed total dose of 15 ml. To obtain the best results accurately assess the body weight.

Administration:

By subcutaneous route. Massage the site after injection to minimize the risk of local reaction.

Withdrawal period:

Meat: 21 days Milk: 3 days

Storage Conditions:

Store below 30° C, Protected from light.
Keep out of reach and sight of children.

MANUFACTURED BY:
NAWAN
LABORATORIES (PVT) LTD.

Animal Health Division
136-138, Sector 15, Korangi Industrial Area,
Karachi-Pakistan.